ACL PATELLAR TENDON AUTOGRAFT RECONSTRUCTION REHABILITATION PROTOCOL

	WEIGHT BEARING	BRACE	ROM	EXERCISES
PHASE I 0-4 weeks	Full in Brace	0-1 week : Locked in full extension for ambulation and sleeping	As tolerated	Heel slides, quad/hamstring sets, patellar mobs, gastroc/soleus stretch SLR w/ brace in full extension
		1-4 weeks: Unlocked for ambulation, remove for sleeping**		until quad strength prevents extension lag
				Side-lying hip/core
PHASE II 4-12 weeks	Full	Discontinue at day 28 if patient has no extension lag	Full	Begin toe raises, closed chain quads, balance exercises, hamstring curls, stationary bike, step-ups, front and side planks; advance hip/core
PHASE III 12-16 weeks	Full	None	Full	Advance closed chain strengthening
				Progress proprioception activities
				Begin stairmaster, elliptical at 8 wks; Running straight ahead at 12 weeks
PHASE IV	Full	None	Full	16 wks: Begin jumping
16-24 weeks				20 wks: Advance to sprinting, backward running, cutting/pivoting/changing direction, initiate plyometric program and sport-specific drills
				22 wks: Advance as tolerated
				FSA completed at 22 wks***
PHASE V > 6 months	Full	None	Full and pain- free	Gradual return to sports participation after completion of FSA
				Maintenance program based on FSA

^{*}Modified with concomitantly performed meniscus repair/transplantation or articular cartilage procedure

^{**}Brace may be removed for sleeping after first post-operative visit (day 7-10)

^{***}Completion of FSA (Functional Sports Assessment) not mandatory, but recommended at approx 22 wks post-op for competitive athletes returning to play after rehab