HTO (HIGH TIBIAL OSTEOTOMY) REHABILITATION PROTOCOL

| | WEIGHT BEARING | BRACE | ROM | EXERCISES |
|-----------------------------------|---|---|---|--|
| PHASE I 0-2 weeks | Heel touch only* | On at all times during day and while sleeping** | 0-90 [°] at home | Calf pumps, quad sets SLR in brace, modalities |
| | | Off for hygiene | | |
| PHASE II 2-6 weeks | Heel touch only | Off at night | Maintain full extension and progress flexion to full | Progress non-weight bearing flexibility, modalities |
| | | Open 0-90 and worn daytime only until 6 wks | | Begin floor-based core and glutes exercises |
| | | | | Advance quad sets, patellar mobs, and SLR |
| PHASE III 6 weeks - 8 weeks | Advance 25% weekly and progress to full with normalized gait pattern | None | Full | Advance closed chain quads, progress balance, core/pelvic and stability work |
| | | | | Begin stationary bike at 6 weeks |
| | | | | Advance SLR, floor-based exercise; hip/core |
| PHASE IV 8-16 weeks | Full | None | Full | Progress flexibility/strengthening, progression of functional balance, core, glutes program |
| | | | | Advance bike, add elliptical at 12 wks as tolerated |
| | | | | Swimming okay at 12 wks |
| PHASE V 16-24 wks | Full | None | Full | Advance Phase IV activity |
| | | | | Progress to functional training, including impact activity after 20 wks when cleared by MD |

*WB status to be confirmed on patient's specific PT Rx

**Brace may be removed for sleeping after first post-operative visit (day 7-10)