

Does ACL Reconstruction Alter Natural History?

A Systematic Literature Review of Long-Term Outcomes

Peter N. Chalmers, MD, Nathan A. Mall, MD, Mario Moric, MS, Seth L. Sherman, MD,
George P. Paletta, MD, Brian J. Cole, MD, MBA, and Bernard R. Bach Jr., MD

Investigation performed at the Departments of Orthopaedic Surgery and Anesthesiology, Rush University Medical Center, Chicago, Illinois, the St. Louis Center for Cartilage Restoration and Repair, Regeneration Orthopedics, St. Louis, and the Department of Orthopaedic Surgery, University of Missouri, Columbia, Missouri

Background: Anterior cruciate ligament (ACL) injury can lead to tibiofemoral instability, decreased functional outcomes, and degenerative joint disease. It is unknown whether ACL reconstruction alters this progression at long-term follow-up.

Methods: A systematic literature review of the long-term results (minimum follow-up, more than ten years) after operative intra-articular reconstruction of ACL injuries and after nonoperative management was performed to compare (1) knee stability on physical examination, (2) functional and patient-based outcomes, (3) the need for further surgical intervention, and (4) radiographic outcomes. After application of selection criteria, forty patient cohorts with a mean of 13.9 ± 3.1 years of postoperative follow-up were identified. Twenty-seven cohorts containing 1585 patients had undergone reconstruction, and thirteen containing 685 patients had been treated nonoperatively.

Results: Comparison of operative and nonoperative cohorts revealed no significant differences in age, sex, body mass index, or rate of initial meniscal injury ($p > 0.05$ for all). Operative cohorts had significantly less need for further surgery (12.4% compared with 24.9% for nonoperative, $p = 0.0176$), less need for subsequent meniscal surgery (13.9% compared with 29.4%, $p = 0.0017$), and less decline in the Tegner score (-1.9 compared with -3.1 , $p = 0.0215$). A difference in pivot-shift test results was observed (25.5% pivot-positive compared with 46.6% for nonoperative) but did not reach significance ($p = 0.09$). No significant differences were seen in outcome scores (Lysholm, International Knee Documentation Committee [IKDC], or final Tegner scores) or the rate of radiographically evident degenerative joint disease ($p > 0.05$ for all).

Conclusions: At a mean of 13.9 ± 3.1 years after injury, the patients who underwent ACL reconstruction had fewer subsequent meniscal injuries, less need for further surgery, and significantly greater improvement in activity level as measured with the Tegner score. There were no significant differences in the Lysholm score, IKDC score, or development of radiographically evident osteoarthritis.

Level of Evidence: Therapeutic Level III. See Instructions for Authors for a complete description of levels of evidence.

Peer Review: This article was reviewed by the Editor-in-Chief and one Deputy Editor, and it underwent blinded review by two or more outside experts. It was also reviewed by an expert in methodology and statistics. The Deputy Editor reviewed each revision of the article, and it underwent a final review by the Editor-in-Chief prior to publication. Final corrections and clarifications occurred during one or more exchanges between the author(s) and copyeditors.

Anterior cruciate ligament (ACL) tears are among the most common injuries in young athletes¹. ACL injury has been associated with tibiofemoral instability, decreased functional outcomes, and meniscal injury²⁻⁷. The damage that leads to these sequelae may occur at the time of the initial event because of associated meniscal injury^{8,9}, chondral injury¹⁰,

subchondral bone impaction^{11,12}, hemarthrosis, and associated ligamentous injuries. Alternately, continued ligamentous deficiency may lead to loss of neuromuscular feedback, altered knee kinematics^{13,14}, increased shear and contact stress, an increased rate of meniscal injury¹⁵⁻¹⁷, and an increased rate of chondral injury¹⁸, each of which could accelerate degenerative changes^{1,19}.

Disclosure: None of the authors received payments or services, either directly or indirectly (i.e., via his or her institution), from a third party in support of any aspect of this work. One or more of the authors, or his or her institution, has had a financial relationship, in the thirty-six months prior to submission of this work, with an entity in the biomedical arena that could be perceived to influence or have the potential to influence what is written in this work. Also, one or more of the authors has had another relationship, or has engaged in another activity, that could be perceived to influence or have the potential to influence what is written in this work. The complete **Disclosures of Potential Conflicts of Interest** submitted by authors are always provided with the online version of the article.

Ligamentous reconstruction may protect against continued instability^{15,20,21}, improve functional outcomes, and decrease rates of degenerative joint disease²².

The largest and highest-quality randomized clinical trial comparing operative and nonoperative treatment suggests that there are “no significant differences in self-reported outcomes at 2 years among the subjects treated with rehabilitation plus early ACL reconstruction, those treated with rehabilitation plus delayed ACL reconstruction, and those treated with rehabilitation alone.”²³ However, considerable deterioration can occur between short and long-term follow-up^{10,24}. The sequelae of continued instability are unlikely to be noted without extended clinical and radiographic monitoring. In addition, the decision not to perform an ACL reconstruction may be guided by a dominant treatment strategy based on identification of the patients as a “coper” with the ability to function normally or nearly normally with an inherently unstable knee. To our knowledge, no long-term randomized clinical trial evidence exists, although long-term outcomes have been reported for numerous non-comparative case series.

The purpose of the present study was to conduct a contemporary systematic review of all published clinical studies with a minimum follow-up of more than ten years after reconstruction and/or nonoperative treatment of ACL-deficient knees, evaluating (1) knee stability on physical examination, (2) functional and patient-based outcomes, (3) the need for further surgical intervention (meniscal surgery and all-cause reoperation), and (4) radiographic outcomes. We hypothesized that reconstruction would lead to greater knee stability on physical examination, improved functional outcomes with decreased rates of later meniscal injury, and a decrease in radiographically evident degenerative joint disease compared with nonoperative therapy.

Materials and Methods

Search Strategy

A systematic literature review was performed in March 2012 with use of the PubMed, Cochrane, BMJ Clinical Evidence, and Embase databases. Search terms used included *anterior cruciate ligament*, *ACL*, *graft*, *reconstruct*, *autograft*, *patella hamstring*, *gracilis*, *semi-*, *tendon*, and *long-term*. Exclusion criteria were a follow-up of less than the ten-year minimum, lack of either physical examination findings or clinical data at the time of final follow-up, a sample size of less than ten, surgery on patients with open physes, outcomes that were not segregated by operative and nonoperative treatment, and a language other than English. In addition, studies that employed ligamentous repair, ligamentous augmentation with a synthetic device, or extra-articular reconstruction or augmentation were excluded. The references of each included article and the table of contents of the last two years of *The Journal of Bone and Joint Surgery* (American and British Volumes); *The American Journal of Sports Medicine*; *Clinical Orthopaedics and Related Research*; *Arthroscopy*; and *Knee Surgery, Sports Traumatology, Arthroscopy* were searched manually for any additional studies. Additionally, the librarian at our institution performed an independent search to ensure that no manuscripts were missed. Study authorship and data were cross-checked to prevent patient data duplication, with longer-term data preferentially included and shorter-term data excluded in such a situation. Exceptions were made for studies reporting on different outcomes in the same patient population. We adhered to the PRISMA (Preferred Reporting Items for Systematic reviews and Meta-Analyses) guidelines²⁵. Studies with Level-I through Level-IV evidence were included.

Data Collection

General demographic data were collected for each study (see Appendix). The following information at the time of final follow-up was collected: need for knee surgery for any cause between the initial treatment and final follow-up, need for further meniscal surgery, number of patients with a side-to-side difference of $>5^\circ$ in range of knee motion, number of patients with an abnormal Lachman test, number of patients with an abnormal pivot-shift test, number of patients with a difference of ≥ 3 mm (the most commonly used threshold in the literature)^{19,26-37} on KT-1000 arthrometer testing (MEDmetric, San Diego, California), mean side-to-side difference on maximum manual testing, mean Lysholm score³⁸, mean pre-injury and final follow-up Tegner activity scores as well as the change between these scores³⁹, percentage of patients returning to their pre-injury level of athletic activity, mean Cincinnati Knee Score⁴⁰, mean Knee Injury and Osteoarthritis Outcome Score (KOOS)⁴¹, mean Short Form (SF)-36 score⁴², mean International Knee Documentation Committee (IKDC) score, and mean overall IKDC grade⁴³. The Lysholm, Tegner, and IKDC scores have been validated in this patient population^{43,44}. Study quality was graded with use of the modified Coleman Methodology Score (CMS)⁴⁵.

Osteoarthritis of the knee at the time of final follow-up was assessed with use of the IKDC system⁴³, the Kellgren and Lawrence system⁴⁶, the Ahlbäck system⁴⁷, the Fairbank system⁴⁸, and the Osteoarthritis Radiographic Severity Index (OARSI)⁴⁹. Radiographic outcomes were then divided (a priori) into those with osteoarthritis as defined previously by Øiestad et al.⁵⁰ (any patient with IKDC grade C or greater) and those without; similar divisions were




Fig. 1
PRISMA diagram showing the identification of the included studies.

made on the basis of a Kellgren and Lawrence grade of ≥ 2 , Ahlbäck grade of ≥ 1 , and Fairbank grade of ≥ 2 .

Statistical Analysis

To avoid excessive bias from a single cohort, analysis was attempted only for those outcomes reported in at least three operative and three nonoperative studies. Studies with both nonoperative and operative arms or with multiple graft type subgroups were included in the systematic review as separate cohorts to reduce the effects of correlated data. If the variance or standard deviation was not given, it was calculated from the standard error, 95% confidence interval (CI), range⁵¹, or p value as available. Dichotomous data such as odds ratios and proportions were re-expressed in the form of the standardized mean difference⁵².

Analyses were performed to pool primary outcome measures among cohorts in the same treatment group. Because of the variability inherent in the use of observational data from a diverse set of studies, the conservative strategy of using random-effects modeling was used for each outcome to estimate and adjust for heterogeneity. This involved use of a mixed-effects meta-regression model (one involving a combination of both random and fixed effects)^{53,54}.

For each pooled outcome measure, the outcome in each cohort was weighted according to the inverse of the sample variance of the outcome in the cohort. After weighting and adjusting for sample size, variability, and sampling error, the overall size of the effect of operative intervention on each outcome variable was estimated with use of a mixed-effects meta-regression model. The treatment effect was estimated with use of a maximum-likelihood estimation (and restricted maximum-likelihood estimation) of the mean difference between the operative and nonoperative groups yielded an estimate of the treatment effect. The treatment effect was standardized according to the pooled variance estimate to calculate the estimated effect size and 95% CI.

If necessary, the sign of the estimated effect size was transposed so that a positive value would indicate a better clinical outcome for the operative group compared with the nonoperative group. All analyses were performed with use of SAS software (version 9.2; SAS Institute, Cary, North Carolina); the mixed-effects meta-regression model analyses were performed with use of the MIXED procedure.

Source of Funding

This study did not receive any external funding.

Results

The initial search revealed 952 potentially relevant abstracts, and twenty-nine remained after application of our study selection algorithm. Thirteen of these reported outcomes of nonoperative therapy^{3-7,30,35,55-60} and twenty-four (twenty-seven cohorts) reported outcomes of operative therapy^{12,19,24,26-37,55-63} (Fig. 1). Two of the studies were randomized clinical trials, two were matched cohort series, seven were prospective cohort series, and eighteen were retrospective cohort series. The twenty-seven operative cohorts had a mean sample size of 58.7 (range, twenty-two to 181), with a total of 1585 included subjects and a mean follow-up of 12.9 years (range, ten to sixteen years) (see Appendix). The ACL reconstruction utilized the patellar tendon in twenty-three (85%) of the operative cohorts, the hamstring tendon in three (11%)^{29,33,62}, and the tibialis anterior tendon in one (4%)²⁶. Although most studies used autograft, allograft was used in some cohorts. Reconstruction took place at a mean




Fig. 2
Summary forest plot showing the mean effect size (with 95% CI) for the comparisons between the operative and nonoperative groups with regard to the Lysholm score, percentage of patients with a positive pivot-shift test, IKDC score, percentage of patients who subsequently developed radiographically evident degenerative joint disease (DJD), percentage of patients requiring further surgery, percentage of patients requiring subsequent meniscal surgery, percentage of patients with preoperative meniscal injury, preoperative Tegner score, Tegner score at the final postoperative follow-up, change in Tegner score between the preoperative and final follow-up time points, age, sex, and BMI.




Fig. 3

Forest plot showing the change in Tegner score between the preoperative and final follow-up time points for each operative (blue) and nonoperative (green) cohort as well as the grand mean for operative and nonoperative treatment. Each horizontal bar shows the weighted standard deviation for the cohort, and the size of the square is weighted according to the sample size and standard deviation. The grand mean for each treatment is shown as a comparably colored vertical line, and its corresponding 95% CI is shown by the horizontal line passing through the diamond. Op = operative cohort, nonOp = nonoperative cohort, BPTB = bone-patellar tendon-bone graft, TA = tibialis anterior or tibialis posterior graft, and HT = hamstring tendon graft.

(and standard deviation) of 20.8 ± 11.3 months after injury. Although most studies used an arthroscopic or mini-open technique, several either involved reconstruction through a medial parapatellar arthroscopy^{55,61} or did not indicate the method of reconstruction⁶⁰. The thirteen nonoperative cohorts had a mean sample size of 52.7 (range, eighteen to ninety-four), with a total of 685 included subjects and a mean follow-up of 16.2 years (range, ten to twenty-seven years). Reporting of each of the outcomes of interest varied among the studies (see Appendix). Preoperatively, the two treatment groups were equivalent with respect to age ($p = 0.7958$), sex ($p = 0.5837$), body mass index (BMI) ($p = 0.8732$), Tegner score ($p = 0.4173$), and the percentage of patients with concomitant meniscal damage ($p = 0.3547$) (Fig. 2).

Assessment with the pivot-shift test revealed a trend toward a difference between the treatment groups ($p = 0.0941$), with 46.6% of nonoperatively treated knees and 25.5% of operatively treated knees being unstable to the pivot (Fig. 2). Because fewer than three nonoperative cohorts had reported results for Lachman testing or KT-1000 testing, no analysis could be performed for these outcomes.

Neither the Lysholm score (mean, 84.4 for nonoperative compared with 88.7 for operative, $p = 0.0837$) nor the IKDC

score (mean, 79.2 for nonoperative compared with 84.5 for operative, $p = 0.2944$) were significantly higher in the operative cohorts (Fig. 2). The operative and nonoperative cohorts showed no significant difference in the Tegner score at the time of final follow-up (4.3 compared with 4.8, $p = 0.1351$), although the operative cohorts did show significantly less decline in the Tegner score relative to the preoperative level (mean, -1.9 compared with -3.1 for nonoperative, $p = 0.0215$) (Figs. 2 and 3). Because fewer than three nonoperative cohorts had reported results for the Cincinnati Knee Score, KOOS, or SF-36 score, no analysis could be performed for these tests.

Further knee surgery was required approximately twice as frequently in the nonoperative cohorts than in the operative cohorts (24.9% compared with 12.4%) (Fig. 4), with the difference in reoperation rate being significant ($p = 0.0176$). The nonoperative cohorts also required subsequent meniscal surgery more than twice as frequently (29.4% compared with 13.9% in the operative cohorts) (Fig. 5), and this difference was also significant ($p = 0.0017$).

Radiographically evident degenerative joint disease was observed in a similar number of patients in the operative and nonoperative cohorts (35.3% compared with 32.8%, $p = 0.7678$) (Fig. 2).




Fig. 4 Forest plot showing the mean percentage of patients requiring further surgery in each operative (blue) and nonoperative (green) cohort for which this outcome was reported as well as the two grand means. See the Figure 3 legend for further details.




Fig. 5 Forest plot showing the percentage of patients requiring subsequent meniscal surgery in each operative (blue) and nonoperative (green) cohort for which this outcome was reported as well as the two grand means. See the Figure 3 legend for further details.

Discussion

ACL deficiency can lead to symptomatic instability, further intra-articular damage, and accelerated degenerative joint disease²⁻⁷. ACL reconstruction is often recommended to protect against continued instability^{20,21}, to improve knee function, to reduce the likelihood of meniscal tears, and possibly to decrease the rate of degenerative joint disease²². However, it is currently unknown whether reconstruction actually confers these benefits. The purpose of this study was to conduct a systematic review of published clinical studies with more than a minimum of ten years of follow-up of reconstructed and nonoperatively treated ACL-deficient knees, evaluating (1) knee stability on physical examination, (2) functional and patient-based outcomes, (3) the need for further surgical intervention (meniscal surgery and all-cause reoperation), and (4) radiographic outcomes.

This study revealed that, compared with nonoperatively treated patients, patients who had undergone ACL reconstruction using modern techniques had fewer subsequent meniscal tears, less need for further surgery, and a greater change in activity from the preoperative level as measured with the Tegner score. No significant differences were observed in the Lysholm score, IKDC score, or radiographic progression of arthritis. Given that the minimum detectable change in the Lysholm score for this population is 8.9⁴⁴ and that the minimum clinically important difference for the IKDC score in a similar population is 16.7⁶⁴, the observed differences in the Lysholm and IKDC scores were also not clinically relevant.

The natural history association between ACL injury, subsequent meniscal injury^{65,66}, subsequent knee surgery, and eventual degenerative joint disease has been described with longitudinal studies¹⁵⁻¹⁷. Our review suggests that patients who undergo ACL reconstruction are less likely to undergo subsequent meniscal surgery compared with patients with ACL deficiency treated nonoperatively. The mean time between ACL injury and ACL reconstruction in this review was 20.8 months, during which time substantial meniscal or articular damage could occur, possibly exacerbating differences in meniscal injury between operative and nonoperative management. This review confirmed that ACL reconstruction protects the knee from the need for further operative interventions. However, the connection between meniscal injury occurring as a result of continued ligamentous instability and the subsequent development of radiographically evident degenerative joint disease remains unclear^{8,9,67}. The lack of any difference in the rate of degenerative joint disease between reconstructed and nonreconstructed knees in our analysis suggests that the continued ligamentous instability and meniscal injury may not be the sole causes of joint degeneration. Further research regarding the role of meniscal injury in the development of degenerative joint disease is necessary⁸⁻¹². Changes in patient activity level may also influence the subsequent development of degenerative joint disease, and a greater increase in activity was noted in operatively treated patients. The development of radiographic and clinical signs of osteoarthritis is likely due to a combination of the effects of the initial trauma⁸⁻¹², meniscal pathology, subtle

rotational instability, patient factors, and biochemical and genetic factors that are incompletely understood^{1,50}. Our study was limited to radiographic evidence of degenerative joint disease, which serves as an imperfect marker of symptomatic degenerative joint disease.

The authors of previous series have also noted that the subjective stability conferred by ACL reconstruction allows patients to resume their activities^{36,37,61,68}, whereas those patients with unreconstructed knees and continued instability often self-limit their activities^{30,35,59}. Our data confirmed this difference, with the reconstructed cohorts experiencing a significantly smaller decrease in the Tegner score compared with the nonreconstructed cohorts. Patients with high expectations with regard to athletic activity should be counseled regarding the change in activity level associated with nonoperative management. Although no significant differences were observed in the Lysholm or IKDC scores, these outcome instruments may be insufficiently sensitive to demonstrate the subtle differences in the quality and stability of an individual knee that prevent a return to higher activity levels^{1,69}.

Previous studies have noted resolution of the pivot-shift phenomenon with reconstruction^{27,70,71}. The present review revealed a 21% difference in the percentage of patients with a positive pivot-shift test in the operative and nonoperative cohorts (26% compared with 47%), but this difference did not reach significance ($p = 0.0941$). The percentage of pivot-positive knees in the operative cohorts was very high compared with the 5% to 9% rates in recent series involving modern ACL reconstruction techniques. However, the recent techniques do not yet have ten-year follow-up data to allow a true comparison, and grafts may stretch between short and long-term follow-up. Alternatively, the high rate of persistence of the pivot phenomenon in the operative cohorts in the present review may reflect nonanatomic tunnel placement in older series^{27,70,71}. Considerable change has occurred in our understanding of the effect of femoral tunnel positioning on rotational tibiofemoral stability during the thirty-year period encompassed by the studies in the present review⁷²⁻⁷⁶. Several trials have demonstrated significantly better rotational stability with anatomic reconstruction⁷⁷⁻⁸⁰; the anatomic femoral tunnel placement may be achievable with transtibial, anteromedial, and two-incision techniques⁸¹⁻⁸³. The present review may thus underestimate the differences between reconstructed and nonreconstructed cohorts. In addition, the pivot maneuver is difficult to perform correctly; subtle variations can alter the results, and thus heterogeneity could have obscured relevant differences among cohorts^{84,85}.

Several previous systematic reviews have been conducted to compare operative and nonoperative treatment of ACL injury^{50,86}. The analysis reported by Linko et al. in 2005 was limited to randomized trials, and the authors did not find sufficient evidence to recommend either reconstruction or rehabilitation alone⁸⁶. In 2009, Øiestad et al. reported the only previous systematic review of outcome studies that was limited to long-term outcomes (at more than ten years)⁵⁰. The authors focused their analysis on radiographically evident degenerative joint disease

and found no difference between the operative and nonoperative cohorts⁵⁰. Of note, over one-half of the studies included in the present review were released since the literature search by Øiestad et al. was performed. Several randomized clinical trials have been conducted to compare operative and nonoperative treatment in the short term⁸⁶; the largest and highest-quality of these suggests that there are “no significant differences in self-reported outcomes at 2 years among the subjects treated with rehabilitation plus early ACL reconstruction, those treated with rehabilitation plus delayed ACL reconstruction, and those treated with rehabilitation alone.”²³ That same study described thirty-two episodes of clinical instability and meniscal signs and symptoms in the delayed reconstruction group compared with three in the early reconstruction group, confirming our findings²³.

The present study has a number of limitations. First, by design, the analysis compared cohort studies performed by different authors. Heterogeneity among these studies limits the interpretation of the results. Considerable changes have occurred between the publication of the first included trial (1993)⁷ and the last (2012)^{29,31} with respect to rehabilitation protocols, patient expectations, and reconstruction techniques. To mitigate this effect, only intra-articular reconstruction techniques were included, with the oldest study of reconstructive outcomes published in 2001. Second, as with any systematic review, the quality of the original data limits the quality of our findings. Third, the radiographic osteoarthritis grade was determined by a single unblinded orthopaedist or radiologist in most of the included studies, limiting the validity of this information. Fourth, limitation of the included data to published studies may have also introduced publication bias. Fifth, the majority of the studies included in this review were published outside the United States (reflecting a bias in the underlying literature), and thus our results may not be generalizable. However, as these five limitations affect both operative and nonoperative data equally, our results are unlikely to have been affected. Sixth, the extended length of time between injury and reconstruction (mean, twenty-one months) may have reduced the potential benefits of reconstruction. Finally, the majority of reconstructions in this review used patellar tendon autograft. Although several studies have noted equivalence between patellar tendon and hamstring tendon autografts^{62,87,88}, it remains unclear whether the results in the present review would also apply to hamstring grafts.

Although no baseline differences between the operative and nonoperative cohorts existed with respect to the percentage of female patients, preoperative Tegner score, or percentage of patients with concomitant meniscal injury, unmeasured residual bias likely exists between the treatment groups. Only

randomization would be able to overcome this limitation; however, most physicians would agree that ACL reconstruction is the treatment of choice in a young, active patient who wishes to return to cutting and pivoting activities^{22,86,89}.

In conclusion, at a mean of 13.9 ± 3.1 years after injury, the patients who underwent ACL reconstruction had fewer subsequent meniscal injuries, less need for further surgery, and a significantly greater improvement in activity level as measured with the Tegner score. Clinical knee stability as measured with the pivot-shift test was also greater, but this difference did not reach significance. There were no significant differences in Lysholm and IKDC scores. Patients who had undergone operative ACL reconstruction did not have a lower rate of radiographically evident osteoarthritis compared with patients who had undergone nonoperative treatment.

Appendix

 Tables summarizing the included studies and the number of studies with reporting of each outcome are available with the online version of this article as a data supplement at jbjs.org. ■

Peter N. Chalmers, MD
Brian J. Cole, MD, MBA
Bernard R. Bach Jr., MD
Department of Orthopaedic Surgery,
Rush University Medical Center,
1611 West Harrison Street, Suite 200,
Chicago, IL 60612

Nathan A. Mall, MD
St. Louis Center for Cartilage Restoration and Repair,
Regeneration Orthopedics,
6 McBride and Sons Center Drive,
Suite 204, St. Louis, MO 63005.
E-mail address: nathanmall@yahoo.com

Mario Moric, MS
Department of Anesthesiology,
Rush University Medical Center,
1653 West Congress Parkway,
Jelke 7, Chicago, IL 60612

Seth L. Sherman, MD
George P. Paletta, MD
Department of Orthopaedic Surgery,
University of Missouri,
1100 Virginia Avenue,
Columbia, MO 65212

References

1. Lohmander LS, Englund PM, Dahl LL, Roos EM. The long-term consequence of anterior cruciate ligament and meniscus injuries: osteoarthritis. *Am J Sports Med.* 2007 Oct;35(10):1756-69. Epub 2007 Aug 29.
2. Sommerlath KG, Gillquist J. The effect of anterior cruciate ligament resection and immediate or delayed implantation of a meniscus prosthesis on knee joint biome-

- chanics and cartilage. An experimental study in rabbits. *Clin Orthop Relat Res.* 1993 Apr;289:267-75.
3. Ageberg E, Pettersson A, Fridén T. 15-year follow-up of neuromuscular function in patients with unilateral nonreconstructed anterior cruciate ligament injury initially treated with rehabilitation and activity modification: a longitudinal

- prospective study. *Am J Sports Med.* 2007 Dec;35(12):2109-17. Epub 2007 Aug 16.
4. Kostogiannis I, Ageberg E, Neuman P, Dahlberg L, Fridén T, Roos H. Activity level and subjective knee function 15 years after anterior cruciate ligament injury: a prospective, longitudinal study of nonreconstructed patients. *Am J Sports Med.* 2007 Jul;35(7):1135-43. Epub 2007 Mar 9.
 5. Messner K, Maletius W. Eighteen- to twenty-five-year follow-up after acute partial anterior cruciate ligament rupture. *Am J Sports Med.* 1999 Jul-Aug;27(4):455-9.
 6. Neuman P, Kostogiannis I, Fridén T, Roos H, Dahlberg LE, Englund M. Knee laxity after complete anterior cruciate ligament tear: a prospective study over 15 years. *Scand J Med Sci Sports.* 2012 Apr;22(2):156-63. Epub 2010 Jul 29.
 7. Neyret P, Donell ST, Dejour H. Results of partial meniscectomy related to the state of the anterior cruciate ligament. Review at 20 to 35 years. *J Bone Joint Surg Br.* 1993 Jan;75(1):36-40.
 8. Englund M, Lohmander LS. Risk factors for symptomatic knee osteoarthritis fifteen to twenty-two years after meniscectomy. *Arthritis Rheum.* 2004 Sep;50(9):2811-9.
 9. Yamaguchi S, Sasho T, Tsuchiya A, Wada Y, Moriya H. Long term results of anterior cruciate ligament reconstruction with iliotibial tract: 6-, 13-, and 24-year longitudinal follow-up. *Knee Surg Sports Traumatol Arthrosc.* 2006 Nov;14(11):1094-100. Epub 2006 Jul 15.
 10. Pernin J, Verdonk P, Si Selmi TA, Massin P, Neyret P. Long-term follow-up of 24.5 years after intra-articular anterior cruciate ligament reconstruction with lateral extra-articular augmentation. *Am J Sports Med.* 2010 Jun;38(6):1094-102. Epub 2010 Mar 19.
 11. Buckwalter JA. Articular cartilage injuries. *Clin Orthop Relat Res.* 2002 Sep;402:21-37.
 12. Hanypsiak BT, Spindler KP, Rothrock CR, Calabrese GJ, Richmond B, Herrenbruck TM, Parker RD. Twelve-year follow-up on anterior cruciate ligament reconstruction: long-term outcomes of prospectively studied osseous and articular injuries. *Am J Sports Med.* 2008 Apr;36(4):671-7. Epub 2008 Mar 7.
 13. Dunn WR, Lyman S, Lincoln AE, Amoroso PJ, Wickiewicz T, Marx RG. The effect of anterior cruciate ligament reconstruction on the risk of knee reinjury. *Am J Sports Med.* 2004 Dec;32(8):1906-14.
 14. Meunier A, Odensten M, Good L. Long-term results after primary repair or non-surgical treatment of anterior cruciate ligament rupture: a randomized study with a 15-year follow-up. *Scand J Med Sci Sports.* 2007 Jun;17(3):230-7.
 15. Cipolla M, Scala A, Gianni E, Puddu G. Different patterns of meniscal tears in acute anterior cruciate ligament (ACL) ruptures and in chronic ACL-deficient knees. Classification, staging and timing of treatment. *Knee Surg Sports Traumatol Arthrosc.* 1995;3(3):130-4.
 16. Mitsou A, Vallianatos P. Meniscal injuries associated with rupture of the anterior cruciate ligament: a retrospective study. *Injury.* 1988 Nov;19(6):429-31.
 17. Nikolić DK. Lateral meniscal tears and their evolution in acute injuries of the anterior cruciate ligament of the knee. *Arthroscopic analysis.* *Knee Surg Sports Traumatol Arthrosc.* 1998;6(1):26-30.
 18. Borchers JR, Kaeding CC, Pedroza AD, Huston LJ, Spindler KP, Wright RW; MOON Consortium and the MARS Group. Intra-articular findings in primary and revision anterior cruciate ligament reconstruction surgery: a comparison of the MOON and MARS study groups. *Am J Sports Med.* 2011 Sep;39(9):1889-93. Epub 2011 Jun 6.
 19. Cohen M, Amaro JT, Ejnisman B, Carvalho RT, Nakano KK, Peccin MS, Teixeira R, Laurino CF, Abdalla RJ. Anterior cruciate ligament reconstruction after 10 to 15 years: association between meniscectomy and osteoarthritis. *Arthroscopy.* 2007 Jun;23(6):629-34.
 20. Bedi A, Musahl V, Steuber V, Kendorf D, Choi D, Allen AA, Pearle AD, Altchek DW. Transtibial versus anteromedial portal reaming in anterior cruciate ligament reconstruction: an anatomic and biomechanical evaluation of surgical technique. *Arthroscopy.* 2011 Mar;27(3):380-90. Epub 2010 Oct 29.
 21. Sim JA, Gadikota HR, Li JS, Li G, Gill TJ. Biomechanical evaluation of knee joint laxities and graft forces after anterior cruciate ligament reconstruction by anteromedial portal, outside-in, and transtibial techniques. *Am J Sports Med.* 2011 Dec;39(12):2604-10. Epub 2011 Sep 9.
 22. Marx RG, Jones EC, Angel M, Wickiewicz TL, Warren RF. Beliefs and attitudes of members of the American Academy of Orthopaedic Surgeons regarding the treatment of anterior cruciate ligament injury. *Arthroscopy.* 2003 Sep;19(7):762-70.
 23. Frobell RB, Roos EM, Roos HP, Ranstam J, Lohmander LS. A randomized trial of treatment for acute anterior cruciate ligament tears. *N Engl J Med.* 2010 Jul 22;363(4):331-42.
 24. Möller E, Weidenhielm L, Werner S. Outcome and knee-related quality of life after anterior cruciate ligament reconstruction: a long-term follow-up. *Knee Surg Sports Traumatol Arthrosc.* 2009 Jul;17(7):786-94. Epub 2009 Apr 10.
 25. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med.* 2009 Jul 21;6(7):e1000097. Epub 2009 Jul 21.
 26. Almqvist KF, Willaert P, De Brabandere S, Criel K, Verdonk R. A long-term study of anterior cruciate ligament allograft reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 2009 Jul;17(7):818-22. Epub 2009 May 7.
 27. Hui C, Salmon LJ, Kok A, Maeno S, Linklater J, Pinczewski LA. Fifteen-year outcome of endoscopic anterior cruciate ligament reconstruction with patellar tendon autograft for "isolated" anterior cruciate ligament tear. *Am J Sports Med.* 2011 Jan;39(1):89-98. Epub 2010 Oct 20.
 28. Lebel B, Hulet C, Galaud B, Burdin G, Locker B, Vielpeau C. Arthroscopic reconstruction of the anterior cruciate ligament using bone-patellar tendon-bone autograft: a minimum 10-year follow-up. *Am J Sports Med.* 2008 Jul;36(7):1275-82. Epub 2008 Mar 19.
 29. Leys T, Salmon L, Waller A, Linklater J, Pinczewski L. Clinical results and risk factors for reinjury 15 years after anterior cruciate ligament reconstruction: a prospective study of hamstring and patellar tendon grafts. *Am J Sports Med.* 2012 Mar;40(3):595-605. Epub 2011 Dec 19.
 30. Meuffels DE, Favejee MM, Vissers MM, Heijboer MP, Reijman M, Verhaar JAN. Ten year follow-up study comparing conservative versus operative treatment of anterior cruciate ligament ruptures. A matched-pair analysis of high level athletes. *Br J Sports Med.* 2009 May;43(5):347-51. Epub 2008 Jul 4.
 31. Murray JRD, Lindh AM, Hogan NA, Trezies AJ, Hutchinson JW, Parish E, Read JW, Cross MV. Does anterior cruciate ligament reconstruction lead to degenerative disease?: Thirteen-year results after bone-patellar tendon-bone autograft. *Am J Sports Med.* 2012 Feb;40(2):404-13. Epub 2011 Nov 23.
 32. Nakata K, Shino K, Horibe S, Tanaka Y, Toritsuka Y, Nakamura N, Koyanagi M, Yoshikawa H. Arthroscopic anterior cruciate ligament reconstruction using fresh-frozen bone plug-free allogeneic tendons: 10-year follow-up. *Arthroscopy.* 2008 Mar;24(3):285-91. Epub 2007 Nov 14.
 33. Sajovic M, Strahovnik A, Dernovsek MZ, Skaza K. Quality of life and clinical outcome comparison of semitendinosus and gracilis tendon versus patellar tendon autografts for anterior cruciate ligament reconstruction: an 11-year follow-up of a randomized controlled trial. *Am J Sports Med.* 2011 Oct;39(10):2161-9. Epub 2011 Jun 28.
 34. Salmon LJ, Russell VJ, Refshauge K, Kader D, Connolly C, Linklater J, Pinczewski LA. Long-term outcome of endoscopic anterior cruciate ligament reconstruction with patellar tendon autograft: minimum 13-year review. *Am J Sports Med.* 2006 May;34(5):721-32. Epub 2006 Jan 6.
 35. Streich NA, Zimmermann D, Bode G, Schmitt H. Reconstructive versus non-reconstructive treatment of anterior cruciate ligament insufficiency. A retrospective matched-pair long-term follow-up. *Int Orthop.* 2011 Apr;35(4):607-13. Epub 2010 Dec 3.
 36. Struwer J, Frangen TM, Ishaque B, Bliemel C, Efe T, Ruchholtz S, Ziring E. Knee function and prevalence of osteoarthritis after isolated anterior cruciate ligament reconstruction using bone-patellar tendon-bone graft: long-term follow-up. *Int Orthop.* 2012 Jan;36(1):171-7. Epub 2011 Sep 7.
 37. van der Hart CP, van den Bekerom MP, Patt TW. The occurrence of osteoarthritis at a minimum of ten years after reconstruction of the anterior cruciate ligament. *J Orthop Surg.* 2008;3:24. Epub 2008 Jun 10.
 38. Lysholm J, Gillquist J. Evaluation of knee ligament surgery results with special emphasis on use of a scoring scale. *Am J Sports Med.* 1982 May-Jun;10(3):150-4.
 39. Tegner Y, Lysholm J. Rating systems in the evaluation of knee ligament injuries. *Clin Orthop Relat Res.* 1985 Sep;198:43-9.
 40. Noyes FR, Barber SD, Moar LA. A rationale for assessing sports activity levels and limitations in knee disorders. *Clin Orthop Relat Res.* 1989 Sep;246:238-49.
 41. Roos EM, Roos HP, Lohmander LS, Ekdahl C, Beynon BD. Knee injury and Osteoarthritis Outcome Score (KOOS)—development of a self-administered outcome measure. *J Orthop Sports Phys Ther.* 1998 Aug;28(2):88-96.
 42. Ware JE Jr, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care.* 1992 Jun;30(6):473-83.
 43. Irrgang JJ, Anderson AF, Boland AL, Harner CD, Kurosaka M, Neyret P, Richmond JC, Shelborne KD. Development and validation of the International Knee Documentation Committee subjective knee form. *Am J Sports Med.* 2001 Sep-Oct;29(5):600-13.
 44. Briggs KK, Lysholm J, Tegner Y, Rodkey WG, Kocher MS, Steadman JR. The reliability, validity, and responsiveness of the Lysholm score and Tegner activity scale for anterior cruciate ligament injuries of the knee: 25 years later. *Am J Sports Med.* 2009 May;37(5):890-7. Epub 2009 Mar 4.
 45. Coleman BD, Khan KM, Maffulli N, Cook JL, Wark JD; Victorian Institute of Sport Tendon Study Group. Studies of surgical outcome after patellar tendinopathy: clinical significance of methodological deficiencies and guidelines for future studies. *Scand J Med Sci Sports.* 2000 Feb;10(1):2-11.
 46. Kellgren JH, Lawrence JS. Radiological assessment of osteo-arthrosis. *Ann Rheum Dis.* 1957 Dec;16(4):494-502.
 47. Ahlback S. Osteoarthritis of the knee. A radiographic investigation. *Acta Radiol Diagn (Stockh).* 1968;Suppl 277:7-72.
 48. Fairbank TJ. Knee joint changes after meniscectomy. *J Bone Joint Surg Br.* 1948 Nov;30(4):664-70.
 49. Altman RD, Hochberg M, Murphy WA Jr, Wolfe F, Lequesne M. Atlas of individual radiographic features in osteoarthritis. *Osteoarthritis Cartilage.* 1995 Sep;3(Suppl A):3-70.

50. Øiestad BE, Engebretsen L, Storheim K, Risberg MA. Knee osteoarthritis after anterior cruciate ligament injury: a systematic review. *Am J Sports Med.* 2009 Jul;37(7):1434-43.
51. Hozo SP, Djulbegovic B, Hozo I. Estimating the mean and variance from the median, range, and the size of a sample. *BMC Med Res Methodol.* 2005;5:13. Epub 2005 Apr 20.
52. Chinn S. A simple method for converting an odds ratio to effect size for use in meta-analysis. *Stat Med.* 2000 Nov 30;19(22):3127-31.
53. Borenstein M, Hedges LV, Higgins JPT, Rothstein HR. *Introduction to meta-analysis.* 1st ed. West Sussex: Wiley; 2009.
54. Hunter JE, Schmidt FL. *Methods of meta-analysis: correcting error and bias in research findings.* 2nd ed. Newbury Park: Sage; 2004.
55. Fink C, Hoser C, Hackl W, Navarro RA, Benedetto KP. Long-term outcome of operative or nonoperative treatment of anterior cruciate ligament rupture—is sports activity a determining variable? *Int J Sports Med.* 2001 May;22(4):304-9.
56. Lohmander LS, Ostergren A, Englund M, Roos H. High prevalence of knee osteoarthritis, pain, and functional limitations in female soccer players twelve years after anterior cruciate ligament injury. *Arthritis Rheum.* 2004 Oct;50(10):3145-52.
57. Mihelic R, Jurdana H, Jotanovic Z, Madjarevic T, Tudor A. Long-term results of anterior cruciate ligament reconstruction: a comparison with non-operative treatment with a follow-up of 17-20 years. *Int Orthop.* 2011 Jul;35(7):1093-7. Epub 2011 Feb 2.
58. Neuman P, Englund M, Kostogiannis I, Fridén T, Roos H, Dahlberg LE. Prevalence of tibiofemoral osteoarthritis 15 years after nonoperative treatment of anterior cruciate ligament injury: a prospective cohort study. *Am J Sports Med.* 2008 Sep;36(9):1717-25. Epub 2008 May 15.
59. Neuman P, Kostogiannis I, Fridén T, Roos H, Dahlberg LE, Englund M. Patellofemoral osteoarthritis 15 years after anterior cruciate ligament injury—a prospective cohort study. *Osteoarthritis Cartilage.* 2009 Mar;17(3):284-90. Epub 2008 Sep 3.
60. von Porat A, Roos EM, Roos H. High prevalence of osteoarthritis 14 years after an anterior cruciate ligament tear in male soccer players: a study of radiographic and patient relevant outcomes. *Ann Rheum Dis.* 2004 Mar;63(3):269-73.
61. Drogset JO, Grøntvedt T, Robak OR, Mølster A, Viset AT, Engebretsen L. A sixteen-year follow-up of three operative techniques for the treatment of acute ruptures of the anterior cruciate ligament. *J Bone Joint Surg Am.* 2006 May;88(5):944-52.
62. Holm I, Øiestad BE, Risberg MA, Aune AK. No difference in knee function or prevalence of osteoarthritis after reconstruction of the anterior cruciate ligament with 4-strand hamstring autograft versus patellar tendon-bone autograft: a randomized study with 10-year follow-up. *Am J Sports Med.* 2010 Mar;38(3):448-54. Epub 2010 Jan 23.
63. Øiestad BE, Holm I, Aune AK, Gunderson R, Myklebust G, Engebretsen L, Fosdahl MA, Risberg MA. Knee function and prevalence of knee osteoarthritis after anterior cruciate ligament reconstruction: a prospective study with 10 to 15 years of follow-up. *Am J Sports Med.* 2010 Nov;38(11):2201-10. Epub 2010 Aug 16.
64. Greco NJ, Anderson AF, Mann BJ, Cole BJ, Farr J, Nissen CW, Irgang JJ. Responsiveness of the International Knee Documentation Committee Subjective Knee Form in comparison to the Western Ontario and McMaster Universities Osteoarthritis Index, modified Cincinnati Knee Rating System, and Short Form 36 in patients with focal articular cartilage defects. *Am J Sports Med.* 2010 May;38(5):891-902. Epub 2009 Dec 31.
65. Dumont GD, Hogue GD, Padalecki JR, Okoro N, Wilson PL. Meniscal and chondral injuries associated with pediatric anterior cruciate ligament tears: relationship of treatment time and patient-specific factors. *Am J Sports Med.* 2012 Sep;40(9):2128-33. Epub 2012 Jun 22.
66. Fok AW, Yau WP. Delay in ACL reconstruction is associated with more severe and painful meniscal and chondral injuries. *Knee Surg Sports Traumatol Arthrosc.* 2013 Apr;21(4):928-33. Epub 2012 May 3.
67. Spindler KP, Huston LJ, Wright RW, Kaeding CC, Marx RG, Amendola A, Parker RD, Andrich JT, Reinke EK, Harrell FE Jr, Dunn WR; MOON Group. The prognosis and predictors of sports function and activity at minimum 6 years after anterior cruciate ligament reconstruction: a population cohort study. *Am J Sports Med.* 2011 Feb;39(2):348-59. Epub 2010 Nov 17.
68. Dunn WR, Spindler KP; MOON Consortium. Predictors of activity level 2 years after anterior cruciate ligament reconstruction (ACLR): a Multicenter Orthopaedic Outcomes Network (MOON) ACLR cohort study. *Am J Sports Med.* 2010 Oct;38(10):2040-50. Epub 2010 Aug 13.
69. Mohtadi N. Development and validation of the quality of life outcome measure (questionnaire) for chronic anterior cruciate ligament deficiency. *Am J Sports Med.* 1998 May-Jun;26(3):350-9.
70. Marcacci M, Zaffagnini S, Giordano G, Iacono F, Presti ML. Anterior cruciate ligament reconstruction associated with extra-articular tenodesis: a prospective clinical and radiographic evaluation with 10- to 13-year follow-up. *Am J Sports Med.* 2009 Apr;37(4):707-14. Epub 2009 Feb 3.
71. Shelbourne KD, Gray T. Minimum 10-year results after anterior cruciate ligament reconstruction: how the loss of normal knee motion compounds other factors related to the development of osteoarthritis after surgery. *Am J Sports Med.* 2009 Mar;37(3):471-80. Epub 2008 Dec 4.
72. Scopp JM, Jasper LE, Belkoff SM, Moorman CT 3rd. The effect of oblique femoral tunnel placement on rotational constraint of the knee reconstructed using patellar tendon autografts. *Arthroscopy.* 2004 Mar;20(3):294-9.
73. Arnold MP, Kooloos J, van Kampen A. Single-incision technique misses the anatomical femoral anterior cruciate ligament insertion: a cadaver study. *Knee Surg Sports Traumatol Arthrosc.* 2001 Jul;9(4):194-9.
74. Giron F, Buzzi R, Aglietti P. Femoral tunnel position in anterior cruciate ligament reconstruction using three techniques. A cadaver study. *Arthroscopy.* 1999 Oct;15(7):750-6.
75. Hantes ME, Zachos VC, Liantsis A, Venouziou A, Karantanas AH, Malizos KN. Differences in graft orientation using the transtibial and anteromedial portal technique in anterior cruciate ligament reconstruction: a magnetic resonance imaging study. *Knee Surg Sports Traumatol Arthrosc.* 2009 Aug;17(8):880-6. Epub 2009 Feb 24.
76. Loh JC, Fukuda Y, Tsuda E, Steadman RJ, Fu FH, Woo SLY. Knee stability and graft function following anterior cruciate ligament reconstruction: comparison between 11 o'clock and 10 o'clock femoral tunnel placement. 2002 Richard O'Connor Award paper. *Arthroscopy.* 2003 Mar;19(3):297-304.
77. Aglietti P, Giron F, Losco M, Cuomo P, Ciardullo A, Mondanelli N. Comparison between single-and double-bundle anterior cruciate ligament reconstruction: a prospective, randomized, single-blinded clinical trial. *Am J Sports Med.* 2010 Jan;38(1):25-34. Epub 2009 Sep 30.
78. Hussein M, van Eck CF, Cretnik A, Dinevski D, Fu FH. Prospective randomized clinical evaluation of conventional single-bundle, anatomic single-bundle, and anatomic double-bundle anterior cruciate ligament reconstruction: 281 cases with 3- to 5-year follow-up. *Am J Sports Med.* 2012 Mar;40(3):512-20. Epub 2011 Nov 15.
79. Ibrahim SA, Hamido F, Al Misfer AK, Mahgoob A, Ghafar SA, Alhran H. Anterior cruciate ligament reconstruction using autologous hamstring double bundle graft compared with single bundle procedures. *J Bone Joint Surg Br.* 2009 Oct;91(10):1310-5.
80. Xu M, Gao S, Zeng C, Han R, Sun J, Li H, Xiong Y, Lei G. Outcomes of anterior cruciate ligament reconstruction using single-bundle versus double-bundle technique: meta-analysis of 19 randomized controlled trials. *Arthroscopy.* 2013 Feb;29(2):357-65.
81. Rue JP, Ghodadra N, Bach BR Jr. Femoral tunnel placement in single-bundle anterior cruciate ligament reconstruction: a cadaveric study relating transtibial lateralized femoral tunnel position to the anteromedial and posterolateral bundle femoral origins of the anterior cruciate ligament. *Am J Sports Med.* 2008 Jan;36(1):73-9.
82. Rue JP, Ghodadra N, Lewis PB, Bach BR Jr. Femoral and tibial tunnel position using a transtibial drilled anterior cruciate ligament reconstruction technique. *J Knee Surg.* 2008 Jul;21(3):246-9.
83. Strauss EJ, Barker JU, McGill K, Cole BJ, Bach BR Jr, Verma NN. Can anatomic femoral tunnel placement be achieving a transtibial technique for hamstring anterior cruciate ligament reconstruction? *Am J Sports Med.* 2011 Jun;39(6):1263-9. Epub 2011 Feb 18.
84. Bach BR Jr, Warren RF, Wickiewicz TL. The pivot shift phenomenon: results and description of a modified clinical test for anterior cruciate ligament insufficiency. *Am J Sports Med.* 1988 Nov-Dec;16(6):571-6.
85. Noyes FR, Grood ES, Cummings JF, Wroble RR. An analysis of the pivot shift phenomenon. The knee motions and subluxations induced by different examiners. *Am J Sports Med.* 1991 Mar-Apr;19(2):148-55.
86. Linko E, Harilainen A, Malmivaara A, Seitsalo S. Surgical versus conservative interventions for anterior cruciate ligament ruptures in adults. *Cochrane Database Syst Rev.* 2005 Apr 18;(2):CD001356.
87. Pinczewski LA, Lyman J, Salmon LJ, Russell VJ, Roe J, Linklater J. A 10-year comparison of anterior cruciate ligament reconstructions with hamstring tendon and patellar tendon autograft: a controlled, prospective trial. *Am J Sports Med.* 2007 Apr;35(4):564-74. Epub 2007 Jan 29.
88. Freedman KB, D'Amato MJ, Nedeff DD, Kaz A, Bach BR Jr. Arthroscopic anterior cruciate ligament reconstruction: a metaanalysis comparing patellar tendon and hamstring tendon autografts. *Am J Sports Med.* 2003 Jan-Feb;31(1):2-11.
89. Andersson C, Odensten M, Gillquist J. Knee function after surgical or nonsurgical treatment of acute rupture of the anterior cruciate ligament: a randomized study with a long-term follow-up period. *Clin Orthop Relat Res.* 1991 Mar;264:255-63.