MPFL RECONSTRUCTION WITH HAMSTRING ALLOGRAFT or MPFL REPAIR/IMBRICATION REHABILIATION PROTOCOL

| | WEIGHT BEARING | BRACE | ROM | EXERCISES |
|-------------------------|----------------------------|--|---|--|
| PHASE I 0-2 weeks | As tolerated with brace | Immobilizer on during day and night Off for hygiene and exercises | 0-90 as tolerated | Quad and hamstring sets, SLR in brace, calf pumps, modalities, heel slides, gastroc stretch |
| PHASE II 2-8 weeks | As tolerated with brace | Begin wearing patellar stabilizing brace after first postop visit. Wear during the day, including exercises Discontinue patellar stabilizing brace at 8 weeks | Advance as tolerated- maintain full extension | Progress weight bearing flexibility, begin toe raises and closed chain quad Begin floor-based core, hip, and glutes, balance work, and hamstring curls Begin stationary bike as tolerated in brace |
| PHASE III 8-16 weeks | Full | None | Full | Advance closed chain quads, progress balance, core, glutes and pelvic stability Begin elliptical, in-line jogging at 12 weeks in PT |
| PHASE IV 4-6 months | Full | None | Full | Progress flexibility and strengthening, progress of functional balance, forward/ backward running, cutting, grapevine, initiate plyometric program and sport-specific drills Return to play after 16 weeks as tolerated and when cleared by MD |

*Brace may be removed at night after first post-operative visit (day 7-14) if directed

**Brace changed to a patellar stabilizing brace (Hely-Weber, Shields, or equivalent) after first postop visit