OSTEOCHONDRAL ALLOGRAFT TO PATELLA/TROCHLEA and FEMORAL CONDYLE REHABILITATION PROTOCOL

	WEIGHT BEARING	BRACE	ROM	EXERCISES
PHASE I 0-8 weeks	**0-6 weeks:** Heel touch only **6-8 weeks:** Advance 25% weekly until full	**0-2 week:** Locked in full extension at all times* Off for CPM, hygiene, and exercises only **2-4 weeks:** Unlocked 0-45°; worn daytime only **4-6 weeks:** Unlocked 0-90°; worn daytime only Discontinue after 6-8 weeks when WB comfortably	**0-2 wks:** Gentle passive 0-45 as tolerated **2 wks+:** advance as tolerated **0-6 wks:** use CPM 6 hours/day, advance 5-10 degrees/day as tolerated, d/c at 6wks Goals: 0-2 wks: 0-30 2-4 wks: 0-60 4-6 wks: 0-90	**0-2 weeks:** Quad sets, SLR, calf pumps, passive leg hangs to 45° at home **2-6 weeks:** PROM/AAROM to tolerance, patella and tibiofibular joint mobs, quad, hamstring, and glute sets/stretching, side-lying hip and core **6-8 weeks:** Addition of heel raises, closed chain exercises, gait normalization, eccentric quads, eccentric hamstrings; advance core, glutes and pelvic stability
PHASE II 8-12 weeks	Full	None	Full	Progress closed chain activities and gait training Begin unilateral stance activities, balance training, hamstring work, hip/core/glutes Begin stationary bike at 10 wks with low resistance as tolerated
PHASE III 12-24 weeks	Full	None	Full	Advance Phase II exercises; maximize core/glutes, pelvic stability work, eccentric hamstrings, balance Swimming as tolerated after 12 weeks Elliptical as tolerated after 16 weeks
PHASE IV 6-12 months	Full	None	Full	Advance functional activity without impact Return to sport-specific activity and impact when cleared by MD after 6-8 months

*Brace may be removed at night after first post-operative visit (day 7-14) if directed

**Alignment correction procedures do not change protocol (ie HTO, DFO, TTO)

BRIAN J. COLE, MD, MBA
MIDWEST ORTHOPAEDICS AT RUSH
BrianColeMD.com

© 2003-2026 Dr. Brian J Cole, MD,MBA. All rights reserved.